

University of Groningen

YidC - an evolutionary conserved device for the assembly of energy-transducing membrane protein complexes

van der Laan, M; Nouwen, NP; Driessen, AJM

Published in:
Current Opinion in Microbiology

DOI:
[10.1016/j.mib.2005.02.004](https://doi.org/10.1016/j.mib.2005.02.004)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2005

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Laan, M., Nouwen, NP., & Driessen, AJM. (2005). YidC - an evolutionary conserved device for the assembly of energy-transducing membrane protein complexes. *Current Opinion in Microbiology*, 8(2), 182-187. <https://doi.org/10.1016/j.mib.2005.02.004>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



ELSEVIER

YidC – an evolutionary conserved device for the assembly of energy-transducing membrane protein complexes

Martin van der Laan¹, Nico P Nouwen² and Arnold JM Driessen²

Members of the YidC/Oxa1/Alb3 membrane protein family are multifunctional mediators of membrane protein integration, folding and assembly into large complexes. Their evolutionary conserved and physiologically important role appears to relate to the assembly of major energy-transducing membrane protein complexes.

Addresses

¹ Institut für Biochemie und Molekularbiologie, Universität Freiburg, Hermann-Herder-Strasse 7, D-79104 Freiburg, Germany

² Molecular Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute and the Materials and Science Centre plus, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands

Corresponding author: Driessen, Arnold JM (a.j.m.driessen@biol.rug.nl)

Current Opinion in Microbiology 2005, **8**:182–187

This review comes from a themed issue on
Cell regulation
Edited by Diego de Mendoza and Ray Dixon

Available online 19th February 2005

1369-5274/\$ – see front matter
© 2005 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.mib.2005.02.004

Introduction

Many mechanistic aspects of intricate biochemical reactions mediated by large membrane protein complexes, such as respiratory electron transport or ion gradient-driven ATP synthesis have been elucidated during the past few years. However, the membrane integration and assembly of these large membrane protein complexes is far from understood.

All primary energy-transducing membranes contain a member of the cytochrome oxidase biogenesis (Oxa) membrane protein family [1,2]. In the inner membrane of mitochondria, Oxa1p mediates the integration of membrane proteins from the matrix side of the membrane. Known substrates of Oxa1p include mitochondrially encoded respiratory chain complex subunits, such as Cox2p, as well as nuclear encoded proteins such as Oxa1 itself [3–5]. In the chloroplast thylakoid membrane, the Alb3 protein is required for the integration of light-harvesting complex proteins (LHCPs) [6]. Bacterial cytoplasmic membranes contain one or more family members that appear to have multiple functions in the biogenesis of membrane proteins. The best-studied example is the YidC protein from the Gram-negative bacterium *Escherichia coli*.

In this review, we discuss the recent insights in the mechanism by which YidC facilitates membrane insertion.

Sec-dependent membrane protein integration

In *E. coli*, most membrane proteins are targeted to the SecYEG translocase via the evolutionary conserved SRP (signal recognition particle) pathway [7]. Translocation of hydrophilic polypeptide domains depends on the ATP-driven motor protein SecA [8–12]. For many membrane proteins the proton motive force (PMF) across the cytoplasmic membrane was shown to be required for insertion and assembly [12–18]. Crosslinking studies have shown that YidC is in close vicinity to transmembrane segments (TMS) of nascent Sec-dependent membrane proteins, such as the cell division protein FtsQ, leader peptidase (Lep) and mannitol permease, during their co-translational insertion [10,11,19–25]. A part of the cellular YidC appears to be physically associated with the Sec translocase [20,26]. Therefore, it was proposed that YidC mediates the lateral release of transmembrane segments from the SecYEG translocase. Several functional studies, however, indicate that YidC is not essential for their membrane insertion. First, *in vivo* depletion of YidC has little effect on the membrane insertion of the monotopic membrane protein FtsQ [10], while its effect on Lep insertion appears to be more pronounced but not critical [21]. Second, a recent *in vitro* study demonstrated that the membrane insertion of FtsQ could be reconstituted with proteoliposomes containing only SecYEG [12]. In this system, the hydrophilic carboxy terminal domain is translocated into the proteoliposome lumen, the TMS of FtsQ was shown to be stably integrated into the lipid bilayer in the absence of YidC. It is still possible that *in vivo* YidC is involved in the hydrophobic partitioning of Sec-dependent TMSs, but at least for FtsQ, it seems that such effects at the most relate to the kinetics of the reaction. Membrane proteins or even their specific TMSs may differ in their YidC requirement, but the features of such TMSs that determine this dependence remain to be elucidated. In addition, YidC seems to play a chaperone role in membrane protein assembly *in vivo*. Experimental support for this hypothesis comes from a recent study by Nagamori *et al.* [27], who have shown that YidC is required for the correct folding of the polytopic membrane protein LacY, while it is dispensable for its membrane integration per se.

Whereas most studies indicate a role of YidC in membrane protein insertion, Fröderberg *et al.* [28] recently

presented evidence for a role of YidC in the Sec-dependent export of lipoproteins. YidC depletion inhibits the translocation of murein lipoprotein (Lpp) and bacteriocin release protein (Brp) *in vivo* and nascent chains of both proteins crosslink to the SRP protein Ffh, SecA, SecY but also to YidC. Since this crosslinking pattern is observed with many nascent membrane proteins, the specific role of YidC in lipoprotein secretion needs to be investigated in more detail.

Sec-independent membrane protein integration

In 2000, Samuelson *et al.* [21] observed that the Sec-independent insertion of the phage protein M13 procoat into the inner membrane of *E. coli* strictly requires YidC *in vivo*. M13 procoat membrane insertion was long believed to be a spontaneous event and a wealth of literature has been published that underscored this hypothesis. M13 procoat consists of two antiparallel TMSs; the first represents a cleavable signal peptide, while the second is a structural unit of the final phage coat. Translocation of the connection loop to the periplasm requires the PMF. As the depletion of YidC was shown to induce strong defects in PMF generation [29•] (see below), this raised the question, whether the effect of YidC depletion on M13 procoat insertion could be indirect. However, PMF-independent point mutants of M13 procoat appear to require YidC for efficient membrane insertion *in vivo* [30] and temperature-sensitive variants of YidC exhibit the M13 procoat insertion defect shortly after a shift to the non-permissive temperature [31], which is earlier than the PMF defects observed upon YidC depletion. This indicates that YidC mediates the membrane insertion of M13 procoat *in vivo*.

Chen *et al.* [32] further demonstrated that a second Sec-independent phage protein, Pf3 coat, requires YidC for membrane insertion both, *in vivo* and *in vitro*. A direct interaction between YidC and Pf3 coat was demonstrated by crosslinking. Pf3 coat is a polypeptide of only 44 amino acids that spans the membrane once with the N-terminus translocated to the periplasm. Therefore, it resembles the second TMS of M13 procoat. Insertion of Pf3 coat is also strongly dependent on the PMF. A recent study by Serek *et al.* [33] showed that the membrane insertion of Pf3 can be reconstituted with proteoliposomes containing only YidC.

What might be the mechanistic role of YidC in Sec-independent membrane protein insertion? For M13 it was shown, that both hydrophobic domains are essential for the translocation of the central loop indicating that an interaction between the two TMSs facilitates membrane insertion [34]. This helical hairpin formation is required because of the high abundance of negative charges in the loop that need to be translocated [35]. Interestingly, although the negative charges acquire a topogenic char-

acter in the presence of a PMF, the synergistic insertion of the two TMSs by hydrophobic interactions alone can drive the translocation [36]. Therefore, it appears that hydrophobic forces and the PMF contribute independently to membrane insertion and that both parameters can at least partly compensate a defect or reduction in the other. For Pf3 coat, a somewhat different scenario seems to exist as the protein contains only one TMS. Mutations in YidC have been identified that specifically affect either M13 procoat or Pf3 coat insertion [30]. The negatively charged residues in the amino terminus of Pf3 coat appear to move electrophoretically across the membrane as this process is driven by the PMF [16]. A mutant Pf3 coat protein (Pf3-4N) that lacks charged residues fails to insert into the membrane [17]. However, when the hydrophobicity of the TMS is increased (Pf3-4N3L), hydrophobic forces drive the spontaneous membrane insertion without topological control [17]. Pf3-4N3L does not require YidC for insertion [33]. These observations confirm the synergistic action of the PMF and hydrophobic forces as described for M13 procoat. However, the interaction of YidC with the Pf3 coat nascent chains is independent of the PMF [32], which supports the notion that YidC and the PMF act on different steps of the insertion reaction.

The data on M13 procoat as well as Pf3 coat strongly suggests that the function of YidC in membrane protein integration is to facilitate the hydrophobic interaction of TMSs with the membrane lipids. It seems likely that YidC provides a special amphiphilic surface that lowers the activation energy of the hydrophobic partitioning event for instance by overcoming the repulsion of the hydrophobic protein segments by polar lipid head groups. In addition, YidC might shield polar residues on their way through the hydrocarbon core of the membrane and promote the folding of hydrophobic regions within a membrane protein into its transmembrane conformation. As in the case of the M13 procoat, this process might involve helical hairpin formation.

Physiological role of YidC

As discussed, YidC probably plays a kinetic role in the process of Sec-dependent membrane protein biogenesis and might act as a chaperone in the organization of helix-helix-interactions without being absolutely required for the insertion of Sec-dependent substrate proteins. In addition, M13 procoat and Pf3 coat strictly depend on YidC for membrane insertion, but these phage proteins do not represent authentic *E. coli* substrates nor are they essential for *E. coli* viability. This raised the question of whether YidC is an essential protein in *E. coli*? A physiological study using a YidC depletion strain showed that YidC depletion causes a strong induction of the membrane stress protein PspA [29•]. PspA induction is correlated with an impairment of the PMF. Further studies revealed that inner membrane vesicles prepared from YidC-depleted cells have a defect to generate a PMF


with both ATP and oxidizable substrates like NADH [29^{••}]. The former correlates with decreased levels of functional F_1F_0 ATP synthase. In particular, the amount of the small, ring-forming rotor subunit c (F_0c) of the membrane-integral F_0 subcomplex is strongly reduced by YidC depletion [29^{••}]. The reduced respiratory capacity is caused by decreased levels of cytochrome *o* oxidase complex from which the amount of the ubiquinol-binding subunit A (CyoA) in the membrane is severely reduced under YidC-depleting conditions [29^{••}]. Since these two enzymes are the key energy-transducing protein complexes of aerobically grown *E. coli*, it seems that the essential role of YidC for *E. coli* viability is related to the biogenesis of these complexes. Yi *et al.* [37] reported that F_1F_0 ATP synthase subunits F_0c as well as F_0 subunit a (F_0a) and the Sec translocase subunit SecE require YidC for membrane insertion *in vivo*. However, F_0a does not stably integrate into the membrane in the absence of F_0c [38], and therefore it is not clear if the insertion defect of F_0a is directly related to the loss of YidC. Also, the observed insertion defect of SecE has to be assessed critically. SecY cannot assemble stably into the membrane in the absence of SecE, and SecE is essential for the functioning of the Sec-translocase. A membrane insertion defect of SecE would imply a strong pleiotropic effect on protein secretion and membrane protein insertion in general. This is not observed in the YidC depletion strain.

The biogenesis of bacterial F_0c was recently studied in detail by van der Laan *et al.* [39^{••}]. F_0c has a molecular mass of 8.3 kDa and consists of two TMSs connected by a small polar loop that is exposed to the cytoplasm. Insertion of *in vitro* synthesized F_0c into YidC-depleted mem-

branes was strongly reduced compared to wild-type membranes. Since F_0c insertion is independent of the PMF, the insertion defect seems directly related to a role of YidC (see Figure 1 for a putative mechanistic model). Importantly, YidC proteoliposomes mediate the efficient and topologically correct membrane insertion of F_0c , while the Sec translocase and the SRP targeting pathway are not required. Although there are conflicting reports on the SRP requirement for F_0c insertion *in vivo* [40,41], the *in vitro* data clearly demonstrate that the SRP pathway is not essential as F_0c insertion into YidC proteoliposomes is not affected by reducing the amounts of either SRP or the SRP receptor FtsY to levels that block the integration of known SRP-dependent substrates, such as FtsQ [39^{••}]. Upon insertion into YidC proteoliposomes, F_0c assembles into large oligomeric complexes in an analogous manner as *in vivo*. These data demonstrate that YidC functions as a membrane protein insertase for the F_0c subunit, which represents the first discovered physiological substrate of this novel insertion pathway. The membrane insertion requirements of F_0c are very similar to that described for M13 and Pf3 procoat (see Figure 2 for a putative mechanistic model). Therefore, it seems that these phage coat proteins employ an essential host cell membrane protein biogenesis pathway to become inserted into the membrane, a prerequisite for phage particle assembly.


Ossenbuhl *et al.* [42^{••}] recently demonstrated that the assembly of the structural core protein D1 into photosystem II in thylakoid membranes of the green alga *Chlamydomonas reinhardtii* is retarded in the absence of Alb3, while membrane integration per se is not affected. In the cyanobacterium *Synechocystis* PC668, deletion of the YidC homolog causes strong defects in the assembly

Figure 1



Model of the YidC-mediated membrane insertion of the F_0 -subunit c. **(a)** F_0 -subunit c initially binds to the membrane via electrostatic interactions. **(b)** Subsequently, YidC provides an amphiphilic surface that facilitates the hydrophobic partitioning of F_0c into the membrane and helical hairpin formation. **(c)** These hydrophobic interactions drive the translocation of the amino- and carboxy-terminal domains across the membrane. Positively charged amino acid residues anchor the hydrophilic loop that connects the two transmembrane segments at the cytosolic surface of the membrane. Next the F_0c subunits assemble into an oligomeric ring structure by a mechanism that is unknown.

Figure 2



Model of the YidC-mediated membrane insertion of M13 procoat. For the insertion of M13 procoat, the common principle of electrostatic membrane binding and hydrophobic partitioning facilitated by YidC is identical as for F_0c insertion (see Figure 1). However, the charge distribution in M13 procoat is inverted. Therefore, the amino- and carboxy-terminal domains are retained in the cytosol while the translocation of the negatively charged hydrophilic loops is strongly stimulated by the proton motive force (PMF). After insertion, procoat is processed by signal peptidase to yield the mature coat that consists of the carboxy-terminal transmembrane segment only.

of the photosystems [43]. These new observations together with the known assembly defects of ATP synthase and cytochrome c oxidase upon deletion of *OXA1* in yeast [44] suggest a general, evolutionary conserved role for Oxa family members in the biogenesis of energy-transducing membrane protein complexes.

Future perspectives

Many questions regarding the mechanistic role of YidC in membrane protein integration and assembly are still open. The mechanistic role of YidC in the Sec-dependent pathway remains unclear. Recently, a direct interaction of Alb3 with the chloroplast SecY has been demonstrated [45]. Mitochondrial inner membranes, however, lack a Sec-type translocase suggesting that only the Sec-independent function of the Oxa family members are evolutionary conserved. In addition, it is not understood how proteins are targeted to the independent YidC/Oxa1/Alb3 insertase. While most studies so far indicate that YidC substrates in bacteria do not require the SRP pathway, a direct interaction of chloroplast SRP/FtsY with Alb3 has been detected [46]. Oxa1 was recently shown to bind mitochondrial ribosomes, which suggest a co-translational insertion mechanism [47,48]. However, the C-terminal domain of Oxa1 required for this interaction is absent in YidC.

The main challenge will be to understand the mechanistic role of YidC/Alb3/Oxa1 in the insertion and assembly of large energy-transducing membrane protein complexes. For instance, how are other membrane-integral subunits of cytochrome *o* oxidase and F_1F_0 ATP synthase in *E. coli* inserted? An initial *in vivo* analysis suggests that subunits a and b of the F_1F_0 ATP synthase depend on the

Sec translocase for membrane integration [40]. The YidC-dependent CyoA protein contains a large, carboxy-terminal periplasmic domain, which likely requires SecYEG and SecA for its correct localization. Therefore, it seems that YidC and SecYEG cooperate in the membrane insertion of CyoA, and this would define CyoA as another class of strictly YidC-dependent proteins next to the F_0c , M13 procoat and Pf3 coat, which require YidC alone.

Taken together, it appears that YidC, Alb3 and Oxa1 have multiple functions in the insertion, folding and assembly of membrane proteins. How they interact with their substrate proteins and how they organize protein–protein interactions within the membrane is currently under investigation.

Acknowledgements

This work was supported by the Earth and Life Sciences Foundation, which is subsidized by the Netherlands Organization for Scientific Research (program grant 809.65.012) and by a fellowship to NN by the Royal Netherlands Academy of Arts and Sciences.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Yen MR, Harley KT, Tseng YH, Saier MH Jr: **Phylogenetic and structural analyses of the oxa1 family of protein translocases.** *FEMS Microbiol Lett* 2001, **204**:223-231.
 2. Kuhn A, Stuart R, Henry R, Dalbey RE: **The Alb3/Oxa1/YidC protein family: membrane-localized chaperones facilitating membrane protein insertion?** *Trends Cell Biol* 2003, **13**:510-516.
 3. He S, Fox TD: **Membrane translocation of mitochondrially coded Cox2p: distinct requirements for export of N and C termini and dependence on the conserved protein Oxa1p.** *Mol Biol Cell* 1997, **8**:1449-1460.

4. Herrmann JM, Neupert W, Stuart RA: **Insertion into the mitochondrial inner membrane of a polytopic protein, the nuclear-encoded Oxa1p.** *EMBO J* 1997, **16**:2217-2226.
5. Hell K, Neupert W, Stuart RA: **Oxa1 acts as a general membrane insertion machinery for proteins encoded by mitochondrial DNA.** *EMBO J* 2001, **20**:1281-1288.
6. Moore M, Harrison MS, Peterson EC, Henry R: **Chloroplast Oxa1p homolog Albino3 is required for post-translational integration of the light harvesting chlorophyll-binding protein into thylakoid membranes.** *J Biol Chem* 2000, **275**:1529-1532.
7. Herskovits AA, Bochkareva ES, Bibi E: **New prospects in studying the bacterial signal recognition particle pathway.** *Mol Microbiol* 2000, **38**:927-939.
8. Lee JI, Kuhn A, Dalbey RE: **Distinct domains of an oligotopic membrane protein are Sec-dependent and Sec-independent for membrane insertion.** *J Biol Chem* 1992, **267**:938-943.
9. Andersson H, von Heijne G: **Sec-dependent and Sec-independent assembly of *E. coli* inner membrane proteins: the topological rules depend on chain length.** *EMBO J* 1993, **12**:683-691.
10. Urbanus ML, Scotti PA, Fröderberg L, Sääf A, de Gier JW, Brunner J, Samuelson JC, Dalbey RE, Oudega B, Lührink J: **Sec-dependent membrane protein insertion: sequential interaction of nascent FtsQ with SecY and YidC.** *EMBO Rep* 2001, **2**:524-529.
11. Urbanus ML, Fröderberg L, Drew D, Björk P, de Gier JW, Brunner J, Oudega B, Lührink J: **Targeting, insertion, and localization of *Escherichia coli* YidC.** *J Biol Chem* 2002, **277**:12718-12723.
12. Van der Laan M, Nouwen N, Driessen AJ: **SecYEG proteoliposomes catalyze the $\Delta\mu$ -dependent membrane insertion of FtsQ.** *J Biol Chem* 2004, **279**:1659-1664.
13. Wolfe PB, Wickner W: **Bacterial leader peptidase, a membrane protein without a leader peptide, uses the same export pathway as pre-secretory proteins.** *Cell* 1984, **36**:1067-1072.
14. Andersson H, von Heijne G: **Membrane protein topology: effects of $\Delta\mu$, H^+ on the translocation of charged residues explain the 'positive inside' rule.** *EMBO J* 1994, **13**:2267-2272.
15. Cao G, Kuhn A, Dalbey RE: **The translocation of negatively charged residues across the membrane is driven by the electrochemical potential: evidence for an electrophoresis-like membrane transfer mechanism.** *EMBO J* 1995, **14**:866-875.
16. Kiefer D, Hu X, Dalbey RE, Kuhn A: **Negatively charged amino acid residues play an active role in orienting the Sec-independent Pf3 coat protein in the *Escherichia coli* inner membrane.** *EMBO J* 1997, **16**:2197-2204.
17. Kiefer D, Kuhn A: **Hydrophobic forces drive spontaneous membrane insertion of the bacteriophage Pf3 coat protein without topological control.** *EMBO J* 1999, **18**:6299-6306.
18. Delgado-Partin VM, Dalbey RE: **The proton motive force, acting on acidic residues, promotes translocation of amino-terminal domains of membrane proteins when the hydrophobicity of the translocation signal is low.** *J Biol Chem* 1998, **273**:9927-9934.
19. Houben EN, Scotti PA, Valent QA, Brunner J, de Gier JL, Oudega B, Lührink J: **Nascent Lep inserts into the *Escherichia coli* inner membrane in the vicinity of YidC, SecY and SecA.** *FEBS Lett* 2000, **476**:229-233.
20. Scotti PA, Urbanus ML, Brunner J, de Gier JW, von Heijne G, van der Does C, Driessen AJ, Oudega B, Lührink J: **YidC, the *Escherichia coli* homologue of mitochondrial Oxa1p, is a component of the Sec translocase.** *EMBO J* 2000, **19**:542-549.
21. Samuelson JC, Chen M, Jiang F, Möller I, Wiedmann M, Kuhn A, Phillips GJ, Dalbey RE: **YidC mediates membrane protein insertion in bacteria.** *Nature* 2000, **406**:637-641.
22. Van der Laan M, Houben EN, Nouwen N, Lührink J, Driessen AJ: **Reconstitution of Sec-dependent membrane protein insertion: nascent FtsQ interacts with YidC in a SecYEG-dependent manner.** *EMBO Rep* 2001, **2**:519-523.
23. Beck K, Eisner G, Trescher D, Dalbey RE, Brunner J, Müller M: **YidC, an assembly site for polytopic *Escherichia coli* membrane proteins located in immediate proximity to the SecYE translocon and lipids.** *EMBO Rep* 2001, **2**:709-714.
24. Houben EN, Urbanus ML, van der Laan M, Ten Hagen-Jongman CM, Driessen AJ, Brunner J, Oudega B, Lührink J: **YidC and SecY mediate membrane insertion of a Type I transmembrane domain.** *J Biol Chem* 2002, **277**:35880-35886.
25. Houben EN, Ten Hagen-Jongman CM, Brunner J, Oudega B, Lührink J: **The two membrane segments of leader peptidase partition one by one into the lipid bilayer via a Sec/YidC interface.** *EMBO Rep* 2004, **5**:970-975.
26. Nouwen N, Driessen AJ: **SecDFyajC forms a heterotetrameric complex with YidC.** *Mol Microbiol* 2002, **44**:1397-1405.
27. Nagamori S, Smirnova IN, Kaback HR: **Role of YidC in folding of polytopic membrane proteins.** *J Cell Biol* 2004, **165**:53-62.
28. Fröderberg L, Houben EN, Baars L, Lührink J, De Gier JW: **Targeting and translocation of two lipoproteins in *Escherichia coli* via the SRP/Sec/YidC pathway.** *J Biol Chem* 2004, **279**:31026-31032.
29. Van der Laan M, Urbanus ML, Ten Hagen-Jongman CM, Nouwen N, Oudega B, Harms N, Driessen A, Lührink J: **A conserved function of YidC in the biogenesis of respiratory chain complexes.** *Proc Natl Acad Sci USA* 2003, **100**:5801-5806.
30. Samuelson JC, Jiang F, Yi L, Chen M, De Gier JW, Kuhn A, Dalbey RE: **Function of YidC for the insertion of M13 procoat protein in *Escherichia coli*: translocation of mutants that show differences in their membrane potential dependence and Sec requirement.** *J Biol Chem* 2001, **276**:34847-34852.
31. Chen M, Xie K, Nouwen N, Driessen AJ, Dalbey RE: **Conditional lethal mutations separate the M13 procoat and Pf3 coat functions of YidC: different YidC structural requirements for membrane protein insertion.** *J Biol Chem* 2003, **278**:23295-23300.
32. Chen M, Samuelson JC, Jiang F, Müller M, Kuhn A, Dalbey RE: **Direct interaction of YidC with the Sec-independent Pf3 coat protein during its membrane protein insertion.** *J Biol Chem* 2002, **277**:7670-7675.
33. Serek J, Bauer-Manz G, Struhalla G, Van Den Berg L, Kiefer D, Dalbey RE, Kuhn A: ***Escherichia coli* YidC is a membrane insertase for Sec-independent proteins.** *EMBO J* 2004, **23**:294-301.
34. Kuhn A, Kreil G, Wickner W: **Both hydrophobic domains of M13 procoat are required to initiate membrane insertion.** *EMBO J* 1986, **5**:3681-3685.
35. Rohrer J, Kuhn A: **The function of a leader peptide in translocating charged amino acyl residues across a membrane.** *Science* 1990, **250**:1418-1421.
36. Kuhn A, Zhu HY, Dalbey RE: **Efficient translocation of positively charged residues of M13 procoat protein across the membrane excludes electrophoresis as the primary force for membrane insertion.** *EMBO J* 1990, **9**:2385-2388.
37. Yi L, Jiang F, Chen M, Cain B, Bolhuis A, Dalbey RE: **YidC is strictly required for membrane insertion of subunits a and c of the F_1F_0 ATP synthase and SecE of the SecYEG translocase.** *Biochemistry* 2003, **42**:10537-10544.
38. Hermolin J, Fillingame RH: **Assembly of F_0 sector of *Escherichia coli* H^+ -ATP synthase.** *J Biol Chem* 1995, **270**:2815-2817.
39. Van der Laan M, Bechtluft P, Kol S, Nouwen N, Driessen AJ: **F_1F_0 ATP synthase subunit c is a substrate of the novel YidC**

pathway for membrane protein biogenesis. *J Cell Biol* 2004, **165**:213-222.

The reconstitution of YidC into proteoliposomes led to the discovery of a novel membrane protein insertion pathway in *E. coli* and the first natural substrate of this pathway. YidC alone mediates the insertion and oligomer assembly of the F_1F_0 ATP synthase rotor subunit c. This process is independent of the Sec translocase and the SRP targeting route.

40. Yi L, Celebi N, Chen M, Dalbey RE: **Sec/SRP requirements and energetics of membrane insertion of subunits a, b, and c of the *Escherichia coli* F_1F_0 ATP synthase.** *J Biol Chem* 2004, **279**:39260-39267.
 41. van Bloois E, Jan Haan G, de Gier JW, Oudega B, Luirink J: **F_1F_0 ATP synthase subunit c is targeted by the SRP to YidC in the *E. coli* inner membrane.** *FEBS Lett* 2004, **576**:97-100.
 42. Ossenbuhl F, Gohre V, Meurer J, Krieger-Liszkav A, Rochaix JD, Eichacker LA: **Efficient assembly of photosystem II in *Chlamydomonas reinhardtii* requires Alb3.1p, a homolog of *Arabidopsis* ALBINO3.** *Plant Cell* 2004, **16**:1790-1800.
- This study shows that chloroplast Alb3 is not required for integration of the plastid-encoded photosystem II core subunit D1 into the thylakoid membrane of *Chlamydomonas reinhardtii*, but the assembly of D1 into functional photosystem II complexes is retarded in an Alb3 mutant. Furthermore, levels of nucleus-encoded light-harvesting proteins are vastly reduced while the remaining antenna systems are still connected to photosystem II reaction centers. These data indicate that Alb3 has a dual function and is required for the assembly of both nucleus- and plastid-encoded protein subunits in photosynthetic complexes of *C. reinhardtii*.
43. Spence E, Bailey S, Nenninger A, Moller SG, Robinson C: **A homolog of Albino3/Oxa1 is essential for thylakoid biogenesis in the cyanobacterium *Synechocystis* PCC6803.** *J Biol Chem* 2004, **279**:5572-5580.
 44. Altamura N, Capitanio N, Bonnefoy N, Papa S, Dujardin G: **The *Saccharomyces cerevisiae* OXA1 gene is required for the correct assembly of cytochrome c oxidase and oligomycin-sensitive ATP synthase.** *FEBS Lett* 1996, **382**:111-115.
 45. Klostermann E, Droste Gen Helling I, Carde JP, Schunemann D: **The thylakoid membrane protein ALB3 associates with the cpSecY-translocase in *Arabidopsis thaliana*.** *Biochem J* 2002, **368**:777-781.
 46. Moore M, Goforth RL, Mori H, Henry R: **Functional interaction of chloroplast SRP/FtsY with the ALB3 translocase in thylakoids: substrate not required.** *J Cell Biol* 2003, **162**:1245-1254.
 47. Jia L, Dienhart M, Schrap M, McCauley M, Hell K, Stuart RA:
 - **Yeast Oxa1 interacts with mitochondrial ribosomes: the importance of the C-terminal region of Oxa1.** *EMBO J* 2003, **22**:6438-6447.

The matrix-localized carboxy terminal domain of Oxa1p binds mitochondrial ribosomes suggesting a co-translational membrane integration mechanism for mitochondrially encoded inner membrane proteins. Truncation of this domain interferes with ribosome binding.
 48. Szyrach G, Ott M, Bonnefoy N, Neupert W, Herrmann JM:
 - **Ribosome binding to the Oxa1 complex facilitates co-translational protein insertion in mitochondria.** *EMBO J* 2003, **22**:6448-6457.

Around the same time as Jia *et al.* [47], this paper also described the binding of mitochondrial ribosomes to the carboxy terminal domain of Oxa1. Furthermore, it demonstrated that the direct ribosome-Oxa1 interaction is crucial for the membrane integration of Cox2, a known Oxa1 substrate.